CLINICAL RESEARCH

Religiousness as a Protective Factor against Psychological Distress in Adult Marijuana Users in the United States

Yorkow Oppon-Acquah^{1*}, Shammah Omololu², Seye Omiyefa³, Victor Ifechukwude Agboli⁴, Lauretta Ekanem Omale⁵, Oluwatosin Kayode⁶, and Adekunle Ade-Onig⁷

¹School of Public and International Affairs, University of Cincinnati, 2800 Clifton Avenue, Clifton Court Hall (CCH), 5th Floor, Cincinnati, OH, USA

²College of Nursing, University of Cincinnati, Procter Hall, 3110 Vine Street, Cincinnati, OH, USA

³Sandra Rosenbaum School of Social Work, University of Wisconsin Madison, 1350 University Ave, Madison, WI

⁴Department of Mathematics and Statistics, Georgia State University, 25 Park Place, 14th Floor, Atlanta, GA, USA

⁵College of Psychology and Behavioral Sciences, National Louis University, 122 S. Michigan Ave., Chicago, IL, USA

⁶The Institute of Design, Illinois Institute of Technology, Kaplan Institute, 3137 South Federal Street, Chicago, IL, USA

⁷Department of Computer Science, Prairie View A & M University, 700 University Drive, Prairie View, TX, USA

Correspondence should be addressed to Yorkow Oppon-Acquah, School of Public and International Affairs, University of Cincinnati, 2800 Clifton Avenue, Clifton Court Hall (CCH), 5th Floor, Cincinnati, OH, USA

Received: 2 May 2024; Accepted: 27 May 2024; Published: 5 June 2024

Copyright © Yorkow Oppon-Acquah. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

SUPPLEMENTAL MATERIAL

Supplemental Table A: Stepwise multiple Poisson regression models estimating effects on psychological distress.

	(1)	(2)	(3)	(4)
	IRR	IRR	IRR	IRR
Variables	(SE)	(SE)	(SE)	(SE)
Religiousness	0.96***	0.96***	0.97***	0.97***
	(0.01)	(0.01)	(0.01)	(0.01)
Marijuana use frequency		1.03***	1.01**	1.01
		(0.01)	(0.00)	(0.00)
Medical marijuana			1.01	1.02*
			(0.01)	(0.012)
Overall health			0.81***	0.81***
			(0.00)	(0.00)
Cigarette use			1.02	1.00
			(0.01)	(0.01)
Binge alcohol use			1.00	0.98
			(0.01)	(0.01)
Illicit drug other than marijuana				1.22***
				(0.01)
Constant	8.78***	7.94***	17.14***	16.81***
	(0.18)	(0.18)	(0.50)	(0.50)
AIC	57318.37	57246.04	54475.03	54207.67

Note: Incidence Rate Ratios (IRR) with standard errors (SE) in parentheses. All models controlled for sociodemographic characteristics which includes age, education, marital status, employment status, sex, sexual orientation, ethnicity/race, and income. The lower value of AIC indicates a better goodness-of-fit of the model. *** p<0.001, ** p<0.01, * p<0.05.